

Blood Feeding Patterns of Potential Arbovirus Vectors of the Genus *Culex* Targeting Ectothermic Hosts

Nathan D. Burkett-Cadena, Sean P. Graham, Hassan K. Hassan, Craig Guyer, Micky D. Eubanks, Charles R. Katholi, and Thomas R. Unnasch*

Department of Entomology and Plant Pathology, and Department of Biological Sciences, Auburn University, Auburn, Alabama; Department of Global Health, University of South Florida, Tampa, Florida; Department of Entomology, Texas A&M University, College Station, Texas; Department of Biostatistics, University of Alabama at Birmingham, Birmingham, Alabama

Abstract. Reptiles and amphibians constitute a significant portion of vertebrate biomass in terrestrial ecosystems and may be important arbovirus reservoirs. To investigate mosquito preference for ectothermic hosts, feeding indices were calculated from data collected in Tuskegee National Forest, Alabama, USA. Four mosquito species fed upon ectothermic hosts, with *Culex peccator* and *Cx. territans* feeding primarily upon ectotherms. These two species appeared to target distinct species with little overlap in host choice. *Culex peccator* was a generalist in its feeding patterns within ectotherms, and *Cx. territans* appeared to be a more specialized feeder. Six of eleven ectotherm species fed upon by *Cx. territans* were fed upon more often than predicted based upon abundance. Spring peepers were highly preferred over other host species by *Cx. territans*. Blood meals taken from each host species varied temporally, with some hosts being targeted fairly evenly throughout the season and others being fed upon in seasonal peaks.

INTRODUCTION

Vector-host interactions are driving forces in the transmission of vector-borne pathogens.¹ An important aspect of vector-host interactions is that of host preference. Each species of hematophagous arthropod feeds on a limited range of host species. Within a group of hosts, however, the degree to which a vector feeds on individual host species can vary tremendously. In addition, even closely related hosts can vary widely in pathogen reservoir competency. Therefore, to accurately describe an arthropod-borne pathogen transmission cycle, it is critical to know which available host species are preferred by the corresponding vectors.

Many studies have reported on the host feeding patterns of mosquitoes.^{2–5} Prior to 1999,⁶ host-preference studies were limited to class-level distinction among mosquito hosts. Although restricted by the available technology, these studies indicated that each species of mosquito fed predominantly on a limited range of hosts (e.g., mammals or birds). Species-level determination of blood meal source, however, suggests that mosquitoes discriminate among hosts beyond the level of class. Hassan and others⁵ reported that ornithophilic mosquitoes feed significantly more or less on available bird species than predicted based on biomass, surface area, or relative abundance. This observation has been confirmed by a number of subsequent studies.^{7–9}

For many of the vector-borne viral encephalitides, the classic transmission scenario involves ornithophilic mosquitoes vectoring virus among avian enzootic hosts in a cycle of amplification.^{10,11} More generalist feeders carry the virus from birds to mammals, enabling the virus to escape the avian enzootic cycle of transmission. Ectotherms, however, may also play a role in this cycle.¹² Reptiles and amphibians make up a large component of vertebrate biomass in terrestrial biological systems,^{13–17} and, somewhat surprisingly, may represent important hosts for several arboviruses. For example,

western equine encephalitis virus may over-winter in garter snakes, *Thamnophis* spp.,^{18,19} and can persist for prolonged periods in the Texas tortoise (*Gopherus berlandieri*).²⁰ Eastern equine encephalitis virus (EEEV) has also been recovered from a number of wild ectotherms,^{21,22} and alligators have been implicated as potential amplifying hosts for West Nile virus.²³ Finally, evidence for the presence of EEEV has been obtained from field-collected ectotherm-feeding mosquitoes, including *Ochlerotatus canadensis*,²⁴ *Uranotaenia sapphirina*,²⁵ *Culex peccator*,¹² and *Culex territans* (Burkett-Cadena ND and others, unpublished data).

In a previous study, we reported that polymerase chain reaction (PCR) assays targeting the cytochrome B gene of vertebrates could be used to identify blood meals derived from ectothermic hosts to the species level¹² in much the same way that such assays have been widely applied to identify avian-derived blood meals. In this study, we have used these PCR-based assays together with abundance data to determine whether ectotherm-feeding mosquitoes are generalists, feeding on available hosts in proportion to host abundance, or if they feed disproportionately on some hosts to a greater extent than would be predicted based upon abundance or biomass alone.

MATERIALS AND METHODS

Study site. Field work was conducted in Tuskegee National Forest, Macon County, Alabama, USA, in an area of the forest with several ponds created by beaver (*Castor canadensis*) activity. All mosquito and ectotherm surveys were conducted within a circle of land (radius = 2 km) with center point on the banks of a beaver pond (32°25.899'N, 85°38.637'W). A detailed description of the site may be found in a previous publication.²⁵

Questing mosquitoes were collected using CDC light traps, as previously described,²⁵ and blooded mosquitoes were collected from natural and artificial resting sites with a battery powered vacuum aspirator. Natural resting sites included cavities in living and dead hardwood trees, herbaceous vegetation, and holes in pond banks created by beavers. Artificial resting sites consisted of a variety of human-made resting shelters, including plastic garbage cans²⁶ and wooden resting

* Address correspondence to Thomas R. Unnasch, Global Health Infectious Disease Research Program, Department of Global Health, College of Public Health, University of South Florida, 3720 Spectrum Boulevard, Suite 304, Tampa, FL 33612. E-mail: tunnasch@health.usf.edu

boxes.²⁷ Collections were begun the first week of February 2007 and continued weekly until October 31, 2007. Blood-engorged females were identified to species²⁸ and stored individually at -70°C for subsequent blood meal identification.

Identification of blood meals. Blood meals were identified to the species level using a modification of a previously described PCR-based assay targeting the vertebrate cytochrome B gene.¹² Briefly, DNA was prepared from individual blooded mosquitoes using the Qiaquick kit (Qiagen, Valencia, CA) following the manufacturer's protocol. A total of 2 μL of the resulting DNA was then used as a template in a 50- μL amplification reaction. The PCR amplifications were conducted in a solution containing 20 mM Tris-HCl, pH 8.4, 50 mM KCl, 2 mM MgCl_2 , 200 μM of each dNTP, 0.5 μM of each primer, and 1.25 units of Taq DNA polymerase (Invitrogen, Carlsbad, CA). Primers used in the PCR were those of Kitano and others²⁹ and were as follows: 5'-GCCTGTT-TACCAAAAACATCAC-3' and 5'-CTCCATAGGGTC-TTCTCGTCTT-3'. Reactions began with incubation at 95°C for 3.5 minutes, followed by 35 cycles at 95°C for 30 seconds, 57°C for 15 seconds, and 72°C for 30 seconds. The reaction was completed by incubation at 72°C for 5 minutes. The PCR products were visualized by electrophoresis on a 1.5% agarose gel and purified by using the QIAquick PCR purification kit (Qiagen). The purified PCR products were subject to direct DNA sequencing, as previously described.³⁰

Estimation of amphibian and reptile abundance. Visual encounter surveys (VESs) were used to estimate relative abundance of amphibians and reptiles at the study site. The VESs are similar to bird point count surveys, and have been widely used in herpetologic research.³¹ Once each week, one diurnal and one nocturnal survey were conducted at five large ponds in the study site. The VESs consisted of an observer(s) walking slowly along the water's edge of a pond and noting all individuals of each species of amphibian or reptile seen. The amount of time spent searching was recorded for each sample period (this varied from 1 to 6 hours depending on conditions), and the relative abundance for each species was calculated as the number of individuals recorded per person-hour of searching.

Because male anurans vocalize, and because recent evidence has suggested that some mosquitoes may use these vocalizations as cues for host choice,³² we also estimated relative abundance of frogs from call surveys.³¹ These surveys were performed during each VES and consisted of recording the identity of each species heard calling. In addition to the identity of each species, the relative abundance of calling males was recorded in one of the following ways: 1) record of exact number of individuals heard calling (1–24); 2) estimate of 25 individuals calling when calls of individuals overlapped but calls of individuals were still distinguishable, and 3) estimate of 50 individuals calling when calls had continuous overlap in which individuals were indistinguishable (see U.S. Geological Survey Patuxent Wildlife Research Center, www.mbr-pwrc.usgs.gov/wifrog/analysis, 2008).

Because amphibians and reptiles vary in size and abundance and because mosquitoes might in part select hosts on factors proportional to based on host biomass,³³ we also collected data to convert our VES estimates of abundance to values of biomass for each species. These values were based on size data from individuals captured during VES samples as

well as opportunistic encounters with individuals at the study site. Each individual was captured by hand and secured in plastic containers for processing. Body mass, measured with a spring scale, was used to calculate biomass by multiplying mean mass of the species by its relative abundance as determined by the VES. Some specimens were transported to the laboratory to obtain blood samples and these animals were weighed on an electronic balance. Blood collections were carried out under procedures approved by the Institutional Animal Care and Usage Committee of Auburn University (protocol no. 2008-1391). For species for which fewer than five mass measurements were obtained, body mass was estimated using the mass of species of similar size.

Feeding index calculation and statistical analysis. Feeding indices were calculated as previously described.⁵ Abundance data used to determine these indices were based on the maximum VES abundance or maximum call index noted for each species during the 2007 field season. Three types of feeding indices were produced. The VES data of ectotherm hosts were used to calculate a visual count feeding index and a biomass-adjusted visual count feeding index. Auditory survey data of anuran hosts were used to calculate a calling-based feeding index. The calling-based index examined whether mosquitoes cue in on vocalizing male anurans regardless of their size and therefore was not adjusted for biomass. Calling-based feeding indexes were calculated for *Cx. territans* only because this species was the only one found to feed upon significant numbers of anurans. All feeding index calculations included only those species actually detected by blood meal analysis. Likelihood ratio tests were used to compare feeding index values, as previously described.⁵ Confidence intervals surrounding the proportion of blood meals obtained from each host class were calculated as previously described.³⁰

RESULTS

A total of 639 blood-engorged female mosquitoes of 13 species were collected during nine months of resting-site aspirations. DNA was extracted from 593 of these samples and subjected to PCR to identify to the species level the source of the blood meals. DNA was extracted from all individual mosquitoes, with the exception that only a random subset of *Anopheles* spp. females were analyzed. Previous studies suggested that mosquitoes of this genus fed exclusively upon mammals.^{3,30} Of the 593 individual mosquitoes from which DNA was extracted, the source of the blood meal was identified for 486 (78%). As expected, *Anopheles crucians*, *An. punctipennis*, and *An. quadrimaculatus* fed exclusively on mammals (Table 1). *Culex quinquefasciatus*, *Culex restuans*, and *Culiseta melanura* fed primarily on birds (Table 1). Four mosquito species (*Cx. territans*, *Cx. peccator*, *Culex erraticus*, and *Cs. melanura*), fed at least some of the time upon ectothermic hosts (Table 1), with *Cx. peccator* and *Cx. territans* feeding primarily upon ectotherms (Table 1). The abundance of non-engorged females of *Cx. territans* peaked in May, and engorged *Cx. territans* exhibited two peaks, in March and in May (Figure 1A). In contrast, populations of both engorged and non-engorged *Cx. peccator* peaked in July (Figure 1B).

In addition to appearing at different times during the season, *Cx. peccator* and *Cx. territans* appeared to partition the ectotherm host community. Of the 14 ectotherm species iden-

TABLE 1
Host class of blood meals identified from field-collected mosquitoes, Tuskegee National Forest, Alabama, 2007

Species	No. meals identified	% Amphibian*	% Avian*	% Mammal*	% Reptile*
<i>Anopheles crucians</i>	10	0	0	100	0
<i>An. punctipennis</i>	7	0	0	100	0
<i>An. quadrimaculatus</i>	8	0	0	100	0
<i>Coquillettidia perturbans</i>	1	0	0	100	0
<i>Culiseta melanura</i>	6	0	83 (53–100)	0	17 (0–47)
<i>Culex erraticus</i>	336	1 (0–2)	23 (19–27)	71 (67–78)	5 (3–7)
<i>Cx. peccator</i>	33	36 (20–52)	3 (0–9)	9 (0–19)	52 (35–69)
<i>Cx. quinquefasciatus</i>	3	0	100	0	0
<i>Cx. restuans</i>	2	0	100	0	0
<i>Cx. territans</i>	78	86 (78–92)	0	4 (0–8)	10 (3–17)
<i>Ochlerotatus sticticus</i>	1	0	0	100	0

* Values are percentages of meals from a given class. Values in parentheses are 95% confidence intervals surrounding that percentage.

tified from blood-fed females, only three host species (the bullfrog [*Rana catesbeiana*], the green frog [*Rana clamitans*], and the southern leopard frog [*Rana sphenoccephala*]) were used by both mosquito species. *Culex peccator* fed primarily upon reptiles, and *Cx. territans* fed primarily upon amphibians. *Culex erraticus* and *Cx. peccator* were the only species that fed on all four classes of vertebrates.

Feeding indices based on VES data suggested that *Cx. peccator* females generally fed on available ectotherm hosts in

proportions that did not significantly differ from those predicted based upon relative abundance, with the exception of the green frog, which was fed upon significantly less ($P = 0.032$) than was predicted based upon its abundance (Figure 2). When host relative abundances were adjusted for biomass, however, the adjusted feeding indices suggest that both the bullfrog ($P = 0.0048$) and the southern leopard frog ($P = 0.0064$) were fed upon in greater proportion than predicted based upon their relative biomass, and cottonmouths were fed upon less frequently ($P = 0.0048$) than predicted by biomass (Figure 2).

Culex territans was more specialized in its host feeding pattern than *Cx. peccator*. Six of 11 ectotherm host species were determined to be fed upon in proportions significantly greater than expected by *Cx. territans* by one or more of the three types of feeding indices (Figure 3). Feeding indices based on VES data suggested that *Cx. territans* females fed on the spring peeper (*Pseudacris crucifer*) ($P = 0.014$) and bullfrogs ($P = 0.016$) in proportions that were significantly greater than that predicted based upon relative abundance alone. When abundance was adjusted for biomass, spring peepers ($P < 0.001$), bird-voiced treefrogs (*Hyla avivoca*) ($P < 0.001$), and squirrel treefrogs (*Hyla squirella*) ($P = 0.047$) were fed upon in proportions greater than predicted. In contrast, feeding indices calculated from calling data indicated that gray treefrogs (*Hyla chrysoscelis*) ($P = 0.031$), bullfrogs ($P = 0.001$), and southern leopard frogs ($P = 0.041$) were fed upon in proportions greater than expected. Two frog species (bullfrogs and spring peepers) were found to be fed upon by *Cx. territans* in proportions significantly greater than predicted in more than one type of feeding index (Figure 3). Five host species (green anole [*Anolis carolinensis*], southern toad [*Bufo terrestris*], green treefrog [*Hyla cinerea*], pine woods treefrog [*Hyla femoralis*], and green frog) were fed upon less than or equal to expected levels, as determined by all three types of feeding indices.

Several abundant species of reptiles and amphibians observed at the study site were not detected in blood meal analysis. For example, cricket frogs (*Acris crepitans* and *A. gryllus*), were by far the most abundant species at the site, representing more than 70% of the total animals observed, and approximately 7% of the total ectotherm biomass (Figure 4). No blood meals, however, were found to have come from these species. Turtles and salamanders, commonly observed throughout the site, were also not detected from blood meals. Spring peepers, in contrast, were the third most commonly




FIGURE 1. Temporal distribution of blood-engorged and non-engorged females of *Culex territans* and *Cx. peccator* from Tuskegee National Forest, Alabama, 2007. Data are normalized for collection events. **A**, Normalized collections of blood engorged and non-engorged *Cx. territans*. **B**, Normalized collections of blood engorged and non-engorged *Cx. peccator*.




FIGURE 2. Feeding indices of ectothermic hosts of *Culex peccator* from Tuskegee National Forest, Alabama, 2007. Feeding indices were calculated as described in the Materials and Methods. In each panel, the feeding indices are shown on a log₂ scale, so that species with feeding indices greater than 1 are indicated as positive bars and those with indices less than 1 as negative bars. Asterisks highlight species where the feeding index value was significantly different than 1.0 ($P < 0.05$). **A**, Feeding indices calculated based upon raw counts from visual encounter surveys of host abundance. **B**, Feeding indices calculated based upon biomass-adjusted visual encounter surveys of host abundance.

fed upon species, yet represented less than 1% of the animals observed and 0.1% of the ectotherm biomass at the site. *Culex territans* fed upon spring peepers 24 times more frequently than predicted, based upon biomass-adjusted abundance (Figure 3B).

Blood meals from ectothermic hosts exhibited considerable temporal variation with respect to host species. Some hosts (such as cottonmouths) were represented in mosquito blood meals fairly evenly throughout the collection season (Figure 5), and other species were unevenly distributed throughout the collection season. Blood meals from spring peepers, for example, were restricted primarily to the late winter and early spring, and blood meals from the gray treefrog were detected in March, April, and May (Figure 5).

DISCUSSION

Previous studies on host preference have suggested that many mosquito species show distinct preferences for certain

host species, such that the feeding pattern cannot be predicted based upon the abundance of the host species alone. This tendency has been demonstrated for mosquitoes that feed on birds^{5,8,9,34} and for mosquitoes that feed on mammals.³⁵ Our own findings indicate that the same is true for some species that feed on reptiles and amphibians. *Culex territans*, which feeds primarily on anurans, was found to be quite host specific, and *Cx. peccator*, which fed on all four classes of terrestrial vertebrates, was less host specific. Given the breadth of hosts from which *Cx. peccator* fed, it is perhaps not surprising that this species showed little apparent preference for one host versus another.

In a previous study conducted at the same site, cotton-




FIGURE 3. Feeding indices of ectothermic hosts of *Culex territans* from Tuskegee National Forest, Alabama, 2007. Feeding indices were calculated as described in the Materials and Methods. In each panel, the feeding indices are shown on a log₂ scale, so that species with feeding indices greater than 1 are indicated as positive bars and those with indices less than 1 as negative bars. Asterisks highlight species where the feeding index value was significantly different than 1.0 ($P < 0.05$). **A**, Feeding indices calculated based upon raw counts from visual encounter surveys of host abundance. **B**, Feeding indices calculated based upon biomass-adjusted visual encounter surveys of host abundance. **C**, Feeding indices calculated based upon calling survey data. NI = not included in analysis (non-vocalizing species).




FIGURE 4. Blood meals, total abundance, and total biomass from ectotherm hosts of *Culex peccator* and *Cx. territans* from Tuskegee National Forest, Alabama, 2007.

mouths and bullfrogs were found to be the host species most commonly fed upon by ectotherm-feeding mosquitoes (primarily *Cx. peccator*).¹² At that time, however, data on the abundance of the various reptiles and amphibians at the site were not available. The current study supports the previous finding that bullfrogs and cottonmouths are commonly targeted hosts. Cottonmouths, however, are very abundant at the Tuskegee National Forest site, representing 41% of the ectotherm biomass. Thus, when considered in the context of their abundance, cottonmouths do not appear to be a preferred host. Bullfrogs, however, despite their relative rarity at the site (total abundance = 0.61% and biomass = 1.85%), appear to be a preferred host for both *Cx. peccator* and *Cx. territans*, with feeding indices greater than 1 in 3 of 5 of the analyses reported.

Spring peepers were another preferred host of *Cx. territans*, as determined by feeding indices calculated from VES data and VES data adjusted for overall biomass. When calling data

were used to calculate feeding indices, however, spring peepers were not found to be a preferred host. In a recent laboratory study, in which *Cx. territans* females were allowed to choose between the calls of multiple frog species, females oriented towards the calls of spring peepers more than any other frog species.³² These findings and the results presented in our report suggest that *Cx. territans* might be specifically attracted to vocalizing spring peepers and other anurans, and may feed actively upon males that are attempting to attract a mate.

In other host groups, e.g., birds, it has been noted that some species are targeted to a lesser extent than would be predicted based upon their abundance. Perhaps the most striking example is that of the American crow, which although being a very common peri-urban bird and one that has been commonly used as a sentinel for West Nile Virus in the United States, appears to be rarely fed upon by the mosquitoes that are the vectors for this virus.^{9,36} Cricket frogs appear to be in a similar position among reptiles and amphibians. These frogs, although abundant at the site, were not detected in mosquito blood meals. Although cricket frogs are (on average) the smallest frogs at the study site, it is unlikely that they are too small to be targeted by mosquitoes. Spring peepers are only slightly larger, and were found to be preferred hosts. It is also unlikely that physical characteristics of the frog's skin (thicker, warty skin in cricket frogs versus thinner, non-warty skin in spring peepers) are responsible for the observed patterns. Although cricket frogs have warty skin (compared with spring peepers), gray treefrogs also have warty skin and




FIGURE 5. Temporal distribution of blood meals taken from preferred ectotherm hosts of *Culex peccator* and *Cx. territans* from Tuskegee National Forest, Alabama, 2007.

were fed upon frequently relative to their abundance. Cricket frogs are sympatric with other frog species at the study site, call during the daytime (as do spring peepers), and have call frequencies similar to that of spring peepers.^{32,37} It is conceivable that cricket frogs are small enough that an approaching mosquito would be considered a prey item and eaten by the frogs. Unlike most hylid frogs (e.g., green treefrog, or spring peeper), cricket frogs apparently continue feeding throughout their extended breeding period.³⁸ Alternatively, it is possible that cricket frogs have skin toxins or other chemical defenses that are repellent to foraging mosquitoes. Amphibians are characterized by a large repertoire of antimicrobial and anti-predatory skin toxins and peptides. The presence or absence of skin toxins has been associated with resistance to fish predation in tadpoles and salamander larvae.³⁹ In addition, some frog skin toxins are associated with resistance to mosquito blood feeding.⁴⁰ Skin semiochemicals associated with mosquito resistance/susceptibility could therefore explain the observed patterns of amphibian host avoidance, particularly in the case of cricket frogs.

Several frog species, including spring peepers, gray treefrogs, and green treefrogs, were fed upon seasonally, and the peak of their appearances in the blood meals coincided with the peak of male calling. This finding indicates that the mosquitoes are finding the frogs at their breeding site, not at the places where they roost for the rest of the year. These temporal patterns provide further support to the hypothesis that *Cx. territans* females are foraging for reproductively active hosts and may locate those hosts by behaviors associated with mate finding. Previous studies have suggested that mosquito feeding upon preferred avian hosts is not evenly distributed temporally, but varies throughout the year. This phenomenon may be a function of the life history or behavior of the host animals^{8,41} and may be an important component of arbovirus amplification. It may also play a role in the process that enables arboviruses to escape avian enzootic cycles and spill over into other vertebrate hosts.⁸ Because ectotherms are potentially important as reservoirs for certain mosquito-borne viruses, it is important to note that similar temporal variation was exhibited for mosquitoes feeding upon ectothermic hosts. That *Cx. peccator* and *Cx. territans* also fed on birds and mammals at our study site points to the possibility that these mosquitoes may be important in the transfer of arboviruses to and from ectotherms and other host groups.

In summary, the data presented above suggest that, like ornithophilic mosquitoes, species targeting ectothermic hosts appear to demonstrate significant biases in host choice. The proportion of blood meals taken from a given host is likely to be a complex function of a number of variables including the innate attractiveness of a species to the mosquito, the availability of that host to the mosquito, and behaviors that make the host either more or less vulnerable to being successfully fed upon by a mosquito. These factors, along with the innate susceptibility of a given vertebrate to a virus are likely to be important in determining the significance of a given species as a reservoir for an arthropod borne virus. In the case of EEEV, a significant amount of evidence has been mounting to support the theory that ectotherms may play a role (of an as yet undetermined magnitude) in the transmission cycle. Our findings that ectotherm feeding mosquitoes prefer some hosts over others could have important consequences for lo-

cal transmission of this pathogen. Additional studies to test these hypotheses are currently underway.

Received May 9, 2008. Accepted for publication July 26, 2008.

Acknowledgments: We thank Nathan Click and Katherine Gray for their help with mosquito collections, Geoffrey Sorrell and Matthew McCurdy for assisting in ectotherm surveys, and David Bayne for graciously allowing access to his land, where a portion of the mosquitoes were collected.

Financial support. This study was supported by a grant from the National Institute of Allergy and Infectious Diseases, Project # R01AI049724 to Thomas R. Unnasch.

Authors' addresses: Nathan Burkett-Cadena, Department of Entomology and Plant Pathology, 301 Funchess Hall, Auburn University, Auburn, AL 36849. Sean Graham and Craig Guyer, Department of Biological Sciences, 331 Funchess Hall, Auburn University, Auburn, AL 36849. Hassan K. Hassan and Thomas R. Unnasch, Global Infectious Disease Research Program, Department of Global Health, College of Public Health, University of South Florida, 3720 Spectrum Boulevard, Suite 304, Tampa, FL 33612. Micky D. Eubanks, Department of Entomology, Texas A&M University, TAMU 2475, College Station, TX 77843-2475. Charles R. Katholi, Department of Biostatistics, University of Alabama at Birmingham, RPHB 327 Birmingham, AL 35294.

REFERENCES

- LoGiudice K, Ostfeld RS, Schmidt KA, Keasing F, 2003. The ecology of infectious disease: effects of host diversity and community composition on Lyme disease risk. *Proc Natl Acad Sci USA* 100: 567–571.
- Edman JD, 1979. Host-feeding patterns of Florida mosquitoes (Diptera: Culicidae) VI. *Culex* (Melanoconion). *J Med Entomol* 15: 521–525.
- Irby WS, Apperson CS, 1988. Hosts of mosquitoes in the coastal plain of North Carolina. *J Med Entomol* 25: 85–93.
- Christensen HA, de Vasquez AM, Boreham MM, 1996. Host-feeding patterns of mosquitoes (Diptera: Culicidae) from central Panama. *Am J Trop Med Hyg* 55: 202–208.
- Hassan HK, Cupp EW, Hill GE, Katholi CR, Klingler K, Unnasch TR, 2003. Avian host preference by vectors of eastern equine encephalomyelitis virus. *Am J Trop Med Hyg* 69: 641–647.
- Boakye D, Tang JM, Truc P, Merriweather A, Unnasch TR, 1999. Identification of blood meals in hematophagous diptera by polymerase chain reaction amplification and heteroduplex analysis. *Med Vet Entomol* 13: 282–287.
- Molaei G, Andreadis TA, Armstrong PM, Anderson JF, Vossbrinck CR, 2006. Host feeding patterns of *Culex* mosquitoes and West Nile virus transmission, northeastern United States. *Emerg Infect Dis* 12: 468–474.
- Kilpatrick AM, Kramer LD, Jones MJ, Marra PP, Daszak P, 2006. West Nile virus epidemics in North America are driven by shifts in mosquito feeding behavior. *PLoS Biol* 4: e82.
- Savage HM, Aggarwal D, Apperson CS, Katholi CR, Gordon E, Hassan HK, Anderson M, Charnetzky D, McMillen L, Unnasch EA, Unnasch TR, 2007. Host choice and West Nile virus infection rates in blood fed mosquitoes, including members of the *Culex pipiens* complex, from Memphis and Shelby County, Tennessee 2002–2003. *Vector Borne Zoonotic Dis* 7: 365–386.
- Shope RE, de Andrade AHP, Bensabath G, Causey OR, Humphrey PS, 1966. The epidemiology of EEE, WEE, SLE and Turlock viruses, with special reference to birds, in a tropical rain forest near Belem, Brazil. *Am J Epidemiol* 84: 467–477.
- Dalrymple JM, Young OP, Eldridge BF, Russell PK, 1972. Ecology of arboviruses in a Maryland freshwater swamp. 3. Vertebrate hosts. *Am J Epidemiol* 96: 129–140.
- Cupp EW, Zhang D, Yue X, Cupp MS, Guyer C, Korves T, Unnasch TR, 2004. Identification of reptilian and amphibian bloodmeals from mosquitoes in an eastern equine encephalomyelitis virus focus in central Alabama. *Am J Trop Med Hyg* 71: 272–276.

13. Burton TM, Likens GE, 1975. Energy flow and nutrient cycling in salamander populations in the Hubbard Brook Experimental Forest, New Hampshire. *Ecology* 56: 1068–1080.
14. Pough FH, 1980. Advantages to ectothermy for tetrapods. *Am Nat* 115: 92–112.
15. Iverson J, 1982. Biomass and turtle populations: a neglected subject. *Oecologia* 55: 69–76.
16. Petranka JW, Murray SS, 2001. Effectiveness of removal sampling for determining salamander density and biomass: a case study in an Appalachian streamside community. *J Herpetol* 35: 36–44.
17. Gibbons JW, Winne CT, Scott DE, Willson JD, Glaudas X, Andrews KM, Todd BD, Fedewa LA, Wilkinson L, Tsaliagos RN, Harper SJ, Greene JL, Tuberville TD, Metts BS, Dorcas ME, Nestor JP, Young CA, Akre T, Reed RN, Buhlmann KA, Norman J, Croshaw DA, Hagen C, Rothermel BB, 2006. Remarkable amphibian biomass and abundance in an isolated wetland: implications for wetland conservation. *Conserv Biol* 20: 1457–1465.
18. Gebhardt LP, Hill DW, 1960. Overwintering of western equine encephalitis virus. *Proc Soc Exp Biol Med* 104: 695–698.
19. Thomas LA, Eklund CM, 1962. Overwintering of western equine encephalomyelitis virus in garter snakes experimentally infected by *Culex tarsalis*. *Proc Soc Exp Biol Med* 109: 421–424.
20. Bowen GS, 1977. Prolonged western equine encephalitis viremia in the Texas tortoise (*Gopherus berlandieri*). *Am J Trop Med Hyg* 26: 171–175.
21. Karstad L, 1961. *Reptiles as Possible Reservoir Hosts for Eastern Encephalitis Virus*. Transactions of the 26th North American Wildlife Conference, 186–202.
22. Hayes RO, Daniels JB, Maxfield HK, Wheeler RE, 1964. Field and laboratory studies on eastern encephalitis in warm- and cold-blooded vertebrates. *Am J Trop Med Hyg* 13: 595–606.
23. Klenk K, Snow J, Morgan K, Bowen R, Stephens M, Foster F, Gordy P, Beckett S, Komar N, Gubler D, Bunning M, 2004. Alligators as West Nile virus amplifiers. *Emerg Inf Dis* 10: 2150–2155.
24. Morris CD, Caines AR, Woodall JP, Bast TF, 1975. Eastern equine encephalomyelitis in upstate New York 1972–1974. *Am J Trop Med Hyg* 24: 986–991.
25. Cupp EW, Klinger K, Hassan HK, Viguers LM, Unnasch TR, 2003. Eastern equine encephalomyelitis virus transmission in central Alabama. *Am J Trop Med Hyg* 68: 495–500.
26. Burkett-Cadena ND, Eubanks MD, Unnasch TR, 2008. Preference of female mosquitoes for natural and artificial resting sites. *J Am Mosq Control Assoc* 24: 228–235.
27. Edman JD, Evans Williams JA, 1968. Development of a diurnal resting box to collect *Culiseta melanura* (Coq). *Am J Trop Med Hyg* 17: 451–456.
28. Darsie RF, Ward RA, 1981. Identification and geographical distribution of the mosquitoes of North America, north of Mexico. *Mosq Systemat* 1S: 1–313.
29. Kitano T, Umetsu K, Tian W, Osawa M, 2007. Two universal primer sets for species identification among vertebrates. *Int J Legal Med* 121: 423–427.
30. Apperson CS, Harrison BA, Unnasch TR, Hassan HK, Irby WS, Savage HM, Aspen SE, Watson DW, Rueda LM, Engber BR, Nasci RS, 2002. Host-feeding habits of *Culex* and other mosquitoes (Diptera: Culicidae) in the borough of Queens in New York City, with characters and techniques for identification of *Culex* mosquitoes. *J Med Entomol* 39: 777–785.
31. Heyer W, Donnelly M, McDiarmid R, Hayek L, Foster M, 1994. *Measuring and Monitoring Biological Diversity. Standard Methods for Amphibians*. Washington, DC: Smithsonian Institution Press.
32. Bartlett-Healy K, Crans W, Gaugler R, 2008. Phonotaxis to amphibian vocalizations in *Culex territans* (Diptera: Culicidae). *Ann Ent Soc Am* 101: 95–103.
33. Khan AA, 1977. Mosquito attractants and repellents. Shorey HH, McKelvey JJ, eds. *Chemical Control of Insect Behavior: Theory and Application*. New York: John Wiley, 305–325.
34. Kilpatrick AM, Daszak P, Jones MJ, Marra PP, Kramer LD, 2006. Host heterogeneity dominates West Nile virus transmission. *Proc R Soc Lond B Biol Sci* 273: 2327–2333.
35. Hess AD, Hayes RO, Tempelis CH, 1968. The use of the foraging ratio technique in mosquito host preference studies. *Mosq News* 28: 386–389.
36. Apperson CS, Hassan HK, Harrison BA, Aspen SE, Savage HM, Farajollahi A, Crans W, Daniels TJ, Falco RC, Benedict M, Anderson M, McMillen L, Unnasch TR, 2004. Host feeding patterns of probable vector mosquitoes of West Nile virus in the eastern United States. *Vector Borne Zoonotic Dis* 4: 71–82.
37. McClelland BE, Wilczynski W, Ryan MJ, 1996. Correlations between call characteristics and morphology in male cricket frogs (*Acris crepitans*). *J Exp Biol* 199: 1907–1919.
38. Wight AH, 1932. *Life-Histories of the Frogs of Okefenokee Swamp, Georgia*. Ithaca, NY: Cornell University Press.
39. Kats C, Petranka J, Sih A, 1988. Antipredator defenses and the persistence of amphibian larvae with fishes. *Ecology* 69: 1865–1870.
40. Williams CR, Smith BP, Best SM, Tyler MJ, 2006. Mosquito repellents in frog skin. *Biol Lett* 2: 242–245.
41. Unnasch RS, Cupp EW, Unnasch TR, 2005. Host selection and its role in transmission of arboviral encephalitides. Collinge SK, Ray C, eds. *Disease Ecology: Community Structure and Pathogen Dynamics*. Oxford, UK: Oxford University Press, 73–89.